

- ▶ **E. REITER**, M. MCNAMARA, J. CLOSSET AND G. HENNEN. Expression and functionality of luteinizing hormone/chorionic gonadotropin receptor in the rat prostate. *Endocrinol.*, 136 : 917-923 (1995).
- ▶ **E. REITER**, O. KECHA, B. HENNUY, S. LARDINOIS, M. KLUG, M. BRUYNINX, J. CLOSSET AND G. HENNEN. Growth Hormone directly affects the function of the different lobes of the rat prostate. *Endocrinol.*, 136 : 3338-3345 (1995).
- ▶ **E. REITER**, B. HENNUY, M. BRUYNINX, A. CORNET, M. KLUG, M. MCNAMARA, J. CLOSSET AND G. HENNEN. Effects of pituitary hormones on the prostate. *Prostate*, 38 : 159-165 (1999).
- ▶ C. TROISPOUX, F. GUILLOU, J.M. ELALOUF, D. FIRSOV, L. IACOVELLI, A. DE BLASI, Y. COMBARNOUS AND **E. REITER**. Involvement of G protein-coupled receptor kinases and arrestins in desensitization to follicle-stimulating hormone action. *Mol. Endocrinol.*, 13 : 1599-1614 (1999).
- ▶ **E. REITER**, S. MARION, F. ROBERT, C. TROISPOUX, F. BOULAY, F. GUILLOU AND P. CRÉPIEUX. Kinase-inactive GRKs are able to attenuate follicle stimulating hormone-induced signaling. *Biophys. Biochem. Res. Comm.*, 282 : 71-78 (2001).
- ▶ P. CRÉPIEUX, S. MARION, N. MARTINAT, V. FAFEUR, Y. LE VERN, D. KERBOEUF, F. GUILLOU AND **E. REITER**. The ERK-dependent signalling is stage-specifically modulated by FSH during primary Sertoli cell maturation. *Oncogene*, 20 : 4696-4709 (2001).
- ▶ S. MARION, F. ROBERT, P. CRÉPIEUX, N. MARTINAT, C. TROISPOUX, F. GUILLOU AND **E. REITER**. GRKs and beta-arrestins are relocalized and attenuate cAMP response to FSH in rat primary Sertoli cells. *Biol. Reprod.*, 66 : 70-76 (2002).
- ▶ P. CRÉPIEUX, C. LECUREUIL, S. MARION, E. KARA, V. PIKETTY, N. MARTINAT, F. GUILLOU, D. ROYERE, **E. REITER**. A mechanistic overview on male infertility and germ cell cancers. *Curr. Pharm. Des.*, 10 : 449-469 (2004).
- ▶ J. KIM, S. AHN, X.R. REN, E.J. WHALEN , **E. REITER**, H. WEI, R.J. LEFKOWITZ. Functional antagonism of different G protein-coupled receptor kinases for beta-arrestin-mediated angiotensin II receptor signaling. *Proc. Nat. Acad. Sci. USA* 102 : 1442-1447 (2005).
- ▶ **E. REITER**, X.R. REN, S. AHN, J. KIM, W. CHEN, R.J. LEFKOWITZ. Different G protein-coupled receptor kinases govern G protein and beta-arrestin-mediated signaling of V2 vasopressin receptor. *Proc. Nat. Acad. Sci. USA* 102 : 1448-1453 (2005).
- ▶ C. LECUREUIL, S. TESSERAUD, E. KARA, N. MARTINAT, A. SOW, I. FONTAINE, C. GAUTHIER, **E. REITER**, F. GUILLOU, P. CRÉPIEUX. Follicle-Stimulating Hormone activates p70 ribosomal protein S6 kinase by protein kinase A-mediated dephosphorylation of Thr 421/ Ser 424 in primary Sertoli cells. *Mol. Endocrinol.* 19 : 1812-1820 (2005).
- ▶ D. GESTY-PALMER, M. CHEN, **E. REITER**, S. AHN, C.D. NELSON, C.R. LOOMIS, R.F. SPURNEY, L.M. LUTTRELL, R.J. LEFKOWITZ. Distinct beta-arrestin and G protein dependent pathways for parathyroid hormone receptor stimulated ERK1/2 activation. *J. Biol. Chem.* 281 : 10856-64 (2006).
- ▶ **E. REITER**, R.J. LEFKOWITZ. 2006 GRKs and beta-arrestins: roles in receptor silencing, trafficking and signaling. *Trends Endocrinol Metab.* 17(4):159-65.
- ▶ S. MARION, E. KARA, P. CRÉPIEUX, V. PIKETTY, N. MARTINAT, F. GUILLOU, **E. REITER**. 2006. G protein-coupled receptor kinase 2 and β -arrestins are re-localized to follicle-stimulating hormone receptor-enriched compartments upon agonist stimulation in rat primary Sertoli cells. *J. Endocrinol.* 190: 343-352.
- ▶ E. KARA, P. CRÉPIEUX, C. GAUTHIER, N. MARTINAT, V. PIKETTY, F. GUILLOU, **E. REITER**. 2006. Identification of a phosphorylation cluster in the C-terminus of the follicle-stimulating hormone receptor (FSH-R) which is important for desensitization and internalization but not β -arrestin-mediated Erk activation. *Mol. Endocrinol.* 20:3014-26.
- ▶ G. BARTHET, B. FRAMERY, F. GAVEN, L. PELLISSIER, **E. REITER**, S. CLAEYSEN, J. BOCKAERT, A. DUMUIS. 2007. 5-HT4 receptors activate ERKs through a pathway independent of G-protein and β -arrestin signalling. *Mol Biol Cell*, 18:1979-91.

- ▶ M. LABASQUE, **E. REITER**, J. BOCKAERT AND P. MARIN. 2008. Physical Interaction of Calmodulin With the 5-HT_{2C} Receptor C-terminus Is Essential for G Protein-Independent, Arrestin-Dependent, Receptor Signaling. *Mol Biol Cell*, 19:4640-50.
- ▶ G. BARTHET, G. CARRAT, E. CASSIER, B. BARKER, F. GAVEN, M. PILLOT, B. FRAMERY, L.P. PELLISSIER, J. AUGIER, D.S. KANG, S. CLAEYSEN, **E. REITER**, J.L. BANÈRES, J.L. BENOVIC, P. MARIN, J. BOCKAERT AND A. DUMUIS. 2009. Beta-arrestin1 phosphorylation by GRK5 regulates G protein-independent 5-HT₄ receptor signalling. *EMBO J.*, 28:2706-18.
- ▶ A. MUSNIER, D. HEITZLER, T. BOULO, S. TESSERAUD, G. DURAND, C. LÉCUREUIL, H. GUILLOU, A. POUPON, **E. REITER**, P. CRÉPIEUX. 2009. Developmental regulation of p70 S6 kinase by a G protein-coupled receptor dynamically modeled in primary cells. *Cell Mol Life Sci*, in press.
- ▶ L. DUPUY, C. GAUTHIER, G. DURAND, A. MUSNIER, D. HEITZLER, A. HERLEDAN, V. SAKANYAN, P. CRÉPIEUX AND **E. REITER**. 2009. A highly sensitive near-infrared fluorescent detection method to analyse signalling pathways by reverse-phase protein array. *Proteomics*, in press.